RESECTABLE AND BORDERLINE RESECTABLE

A. Treatment Naïve
   I. CA19-9 Producer (CA 19-9 > 35 U/mL, when total bilirubin 2 mg/dL)
      a. PANC Trial - Phase II trial of adaptive neoadjuvant therapy
C. Post Surgical Resection
   I. **PROTECT-PANC - Phase II**
   cancer recurrence (NEW!!)
   II. ELICIO-ELI-002-201- Phase
   KRAS/NRAS mutated solid t

LOCALLY ADVANCED

A. Type A – potentially operable
   I. SOFT Trial - Phase II, RCT of IMRT vs SBRT prior to surgery
B. Post Surgical Resection
   I. **PROTECT-PANC - Phase II, adjuvant therapy for patients at risk of cancer recurrence (NEW!!)**
   II. ELICIO-ELI-002-201- Phase I/II, study of ELI-002 7P in subjects with KRAS/NRAS mutated solid tumors

METASTATIC

A. Phase 1 and 2
   I. ASTELLAS-2138-CL-0101- Phase I/Ib, ASP2138 in Adults with Stomach Cancer or Pancreatic Cancer *(check for slot availability)*
   II. HELIX-LDOS006- Phase I/II, Study of the Microenvironment Modifier L-DOS47 Plus Doxorubicin *(check for slot availability)*
   III. MIRATI 1719-001 - A Phase I/II Multiple Expansion Cohort Trial of MRTX1719 in Patients with Advanced Solid Tumors with Homozygous MTAP Deletion *(check for slot availability)*
   IV. IIT-GEORGE-I-PREDICT- Phase I, Investigation of Profile-Related Evidence Determining Individualized Cancer Therapy for Patients

PANCREATIC NEUROENDOCRINE

   I. ALLIANCE-A022001-PNETS- Lutetium LU 177 Dotatate PRRT vs Capecitabine and Temozolomide in PNET
   II. AADI-NET-202: Phase II, IV Nab-Sirolimus in patients with well-

RESECTABLE & BORDERLINE RESECTABLE

Clinical Trial Name: Adaptive Modification of Neoadjuvant Therapy Based on Clinical Response in Patients with Localized Pancreatic Cancer (PANC Trial)

Research Coordinator: Megan Graham **Phone:** 414-805-8921	ECOG performance status < 2 Histologically confirmed adenocarcinoma. Biopsy must have been completed prior to start of treatment Clinical stage consistent with resectable or borderline resectable adenocarcinoma of the pancreas, based on CT or MRI findings $GHTXDWH RUJDQ DQG ERQH PDUURZ IXQFWLRQ DV GHILQHG E\ WRWDO OHXNR SODWHOHWV ! [H / FUHDWLQLQH FOHDUDQFH ! P/ PLQ*R U6 7 HD7L Q L Q H ALT/SGPT <3 x ULN CA19-9 producer, as defined by a pretreatment CA 19-9 > 35 U/mL, when total bilirubin ≤2 mg/dL. Key Exclusion: Received chemotherapy and/or radiation within three years prior to study enrollment Previous history of another malignancy w/in 3 years of study (other than cured basal or squamous cell carcinoma and other in situ carcinomas that were completely treated or localized prostate cancer with normal prostate specific antigen)

Clinical Trial Name: Stereotactic Body Radiation Therapy or Conventionally Fractionated Concurrent Chemotherapy and Radiation Therapy Preoperatively for Resectable or Borderline Resectable Pancreatic Adenocarcinoma (SOFT Trial)

Study Design: This study is a prospective, open-label, randomized, parallel, two-arm, phase II clinical trial. Patients meeting the eligibility criteria will be randomized after a minimum of two months of induction chemotherapy. These patients will be required to have no biopsy-proven distant disease on repeat staging studies before randomization. Patients who have radiologically equivocal evidence of distant metastatic disease (small lung nodules, or liver lesions that cannot be definitively characterized, etc.) are also eligible for enrollment. Patients with biopsy-proven metastatic disease are not eligible.

NCT#: NCT03704662

Key Inclusion

Confirmed, resectable/borderline resectable, locally advanced Type A pancreatic adenocarcinoma
Patients with and without regional adenopathy are eligible
No evidence of distant metastatic disease

NCT#: NCT06048484 **Study PI:** Dr. Ben George **Research Coordinator:** Grace Westerman **Phone:** 414-805-8986	Key Inclusion : Histological or pathological confirmation of pancreatic adenocarcinoma Cytologic or histologic proof of pancreatic ductal adenocarcinoma (PDAC) needs to be verified by the treating institution pathologist. A pathological report from non-treating institutions is sufficient to consent and to initiate investigational therapy if tissue sample is unavailable for evaluation at time of consent or enrollment. However, in such a case, PDAC diagnosis should be confirmed by the treating institution pathologist at a later time. Completed 8 cycles of neoadjuvant modified FOLFIRINOX. Omission of oxaliplatin due to adverse events may be allowed in cycles 5-8 with consultation with the principal investigator. Patients with surgically resectable PDAC who are considered appropriate to undergo the applicable operation. MCW's criteria of borderline resectable meets this inclusion criteria. Eligible to undergo SBRT. Measurable disease as per Response Evaluation Criteria in Solid Tumors (RECIST) 1.1. No prior surgical, systemic, or radiotherapy for PDAC except for mFOLFIRINOX. ECOG: 0 or 1. Key Exclusion : Prior treatment with T-cell co-stimulating or immune checkpoint blockade therapies, including but not limited to anti-CTLA-4, anti-PD-1, and anti-PD-L1 therapeutic antibodies. Uncontrolled pleural effusion, pericardial effusion, or ascites. Uncontrolled hypercalcemia (ionized calcium > 1.5 mmol/L, calcium > 12 mg/dL, or corrected serum calcium > ULN) or symptomatic hypercalcemia requiring continued use of bisphosphonate therapy. Active or history of autoimmune disease or immune deficiency, including, but not limited to, myasthenia gravis, myositis, autoimmune hepatitis, systemic lupus erythematosus, rheumatoid arthritis, inflammatory bowel disease (Crohn's disease or ulcerative colitis), antiphospholipid antibody syndrome, Wegener granulomatosis, Sjögren's syndrome, Guillain-Barré syndrome, or multiple sclerosis (some exceptions permissible as outlined per protocol). History of idiopathic pulmonary fibrosis, interstitial lung disease, organizing pneumonia (e.g., bronchiolitis obliterans), drug-induced pneumonitis, or idiopathic pneumonitis, or evidence of active pneumonitis on screening chest CT scan. History of radiation pneumonitis in the radiation field (fibrosis) is permitted. Positive HIV test at screening or at any time prior to screening. Active hepatitis B virus (HBV) infection (chronic or acute), defined as having a positive hepatitis B surface antigen (HBsAg) test at screening. Active hepatitis C virus (HCV) infection, defined as having a positive HCV antibody test followed by a positive HCV RNA test at screening. The HCV RNA test will be performed only for patients who have a positive HCV antibody test. History of allergy or hypersensitivity to oxaliplatin, irinotecan, leucovorin, fluorouracil, pegfilgrastim, or any excipients. History of Gilbert's disease or known genotype UGT1A1 *28/*28.

Clinical Trial Name: Molecular Profile-related Individualized Targeted Therapy in Resected Pancreatic Cancer with High-Risk of Cancer Recurrence (PROTECT-PANC)

NEW PATIENT COORDINATOR: (414) 805-6849

LOCALLY ADVANCED

Clinical Trial Name: Stereotactic Body Radiation Therapy or Conventionally Fractionated Concurrent Chemotherapy and Radiation Therapy Preoperatively for Resectable or Borderline Resectable Pancreatic Adenocarcinoma (SOFT Trial)

Study Design: This study is a prospective, open-label, randomized, parallel, two-arm, phase II clinical trial. Patients meeting the eligibility criteria will be randomized after a minimum of two months of induction chemotherapy. These patients will be required to have no biopsy-proven distant disease on repeat staging studies before randomization. Patients who have radiologically equivocal evidence of distant metastatic disease (small lung nodules, or liver lesions that cannot be definitively characterized, etc.) are also eligible for enrollment. Patients with biopsy-proven metastatic disease are not eligible.

NCT#: NCT03704662

Study PI: Dr. William Hall

Research Coordinator: Kathryn Hallada

Phone: 414-805-0124

Key Inclusion :

Confirmed, resectable/borderline resectable, locally advanced Type A pancreatic adenocarcinoma
Patients with and without regional adenopathy are eligible
No evidence of distant metastatic disease
F\FOH RI V\VWHPLF FKHPRWKHUDS\ ZLWKRXW HYLGHQFH RI GLVWDQW SURJUH

Key Exclusion :

Distant metastatic disease
Prior invasive malignancy within the last 3 years
Prior radiotherapy to the region of the study cancer that would result in overlap of radiation therapy fields
Major surgery within 28 days prior to study entry

NEW PATIENT COORDINATOR:

NEW PATIENT COORDINATOR:

Active autoimmune disease that has required systemic immunosuppressive treatment within the past 1 month prior to the start of study intervention.

Major surgical procedure 28 days before start of study intervention and has not fully recovered.

Received radiotherapy for locally advanced unresectable or metastatic gastric or GEJ or metastatic pancreatic adenocarcinoma 14 days prior to start of study intervention and has NOT recovered from any related toxicity.

Received an CLDN18.2-targeted investigational agent (e.g., zolbetuximab or chimeric antigen receptor CLDN18.2-specific T cells) prior to first dose of study intervention administration is not eligible for dose escalation cohorts. However, a participant who has

Clinical Trial Name: MRTX1719 in Patients With Advanced Solid Tumors With Homozygous MTAP Deletion (MIRATI 1719-001)

Study Design: The study is a Phase 1/2, open-label, multicenter, study of the safety, tolerability, PK, PD, and anti-tumor activity of MRTX1719 patients with advanced, unresectable or metastatic solid tumor malignancy with homozygous deletion of the MTAP gene.

NCT #:

NCT05245500

Study PI:

Dr. Ben George

Research Coordinator:

Nicholas Pucek

Phone:

414-805-4639

Key Inclusion :

- Histologically confirmed diagnosis of a solid tumor malignancy with homozygous deletion of the MTAP gene detected in tumor tissue or ctDNA
- Unresectable or metastatic disease
- Patients must have received standard therapies appropriate for their tumor type and stage with disease progression on or after the most recent treatment
  - *f*Phase 1 dose escalation, RECIST 1.1 measurable or evaluable disease
  - *f*Phase 1b and Phase 2 cohorts, RECIST 1.1 measurable disease
- Presence of a tumor lesion amenable to mandatory biopsy for pharmacodynamic evaluation at baseline and on-study unless Sponsor-confirmed as medically unsafe or infeasible
- ECOG: 0 or 1

Key Exclusion:

- Prior treatment with a PRMT5 or MAT2A inhibitor therapy.
- Active brain metastases or carcinomatous meningitis.
- History of significant hemoptysis or hemorrhage within 4 weeks of the first dose of study treatment.
- Major surgery within 4 weeks of first dose of study treatment.
- History of intestinal disease, inflammatory bowel disease, major gastric surgery, or other gastrointestinal conditions (eg, uncontrolled nausea, vomiting, malabsorption syndrome) likely to alter absorption of study treatment or result in inability to swallow oral medications
- Cardiac abnormalities

1 TfD-012 0 T39 C0.949 0.949 0.976 rg48 9.96 269.28 11aP 32.7698 Tz (0 0.49 0.949 0.976 rg45.22 323.06 102.48.29 11a3.3349 g-004c 0.002 45.12

Research Coordinator:

Paola Gonzalez Quevedo

Phone:

NEW PATIENT COORDINATOR:

PANCREATIC NEUROENDOCRINE

Clinical Trial Name: ALLIANCE-A022001-PNETS Lutetium LU 177 Dotatate PRRT vs Capecitabine and Temozolomide in PNET

Study Design: This is a phase II randomized, prospective trial of Lutetium LU 177 Dotatate PRRT versus Capecitabine and Temozolomide in well-differentiated pancreatic neuroendocrine tumors.

NCT#:
NCT05247905

Key Inclusion

Histologic or pathologic documentation of well-differentiated pancreatic neuroendocrine tumor (G1, G2, or well-differentiated G3) confirmed by local histology and/or pathology. Functional or nonfunctional tumors are allowed.

Stage: locally unresectable or metastatic disease.

Tumor Site: neuroendocrine tumor of pancreatic primary site.

Radiologic evaluation: tumor must have shown somatostatin receptor (SSTR) positivity on 68Ga-DOTATATE PET or other SSTR- 3 (7 V F D Q L Q W K H P R Q W K V S U L R U W R U H J L V W U D W L R Q K R Z H Y H U G R F X P H Q W